What have we learned?

Pat Brandes USFWS

What have we learned?

- Survival in 2006 appears to have been higher with lower exports
 - No clear relationship between F/E ratio and smolt survival in range tested w/o barrier
 - Survival has been low since 2003
 - Need to measure survival at exports of 1500 with flows of 7000 and HORB
 - Need to continue measuring survival with and without HORB
 - Need to identify sources/locations of mortality


EWA and B2 assets were used in 2006 to reduce exports for high and low export experiment

WHY VAMP?


To better understand how flow/exports affect juvenile salmon survival through the Delta for salmon originating from the SJ tributaries with and without the HORB in place

Conceptual model of how smolt survival may vary with flow/exports

Flow/export vs adult escapement



Mean Flow/Export Ratio between April 15 and June 15 (1951 to 2003) and escapement 2 ½ years later



VAMP release sites in the Sacramento-San Joaquin Estuary, California.


Methods:

- Estimate combined recovery rate
 - from recoveries at Antioch, Chipps Island and in the ocean fishery (as available)
 (no ocean recoveries in 2004-2006)

 Estimate ratio of recovery rates (combined differential recovery rate -CDRR)

Ratio: Recovery Rate of upstream group

Recovery Rate of downstream group




CDRR (+/- 2 SE) using Chipps Island and Antioch recoveries of smolts released at Mossdale (MD) relative to those released at Jersey Point for the first (1), second (2) release groups and their difference in 2006.


Flow

Exports

What have we learned: Survival in 2006 appears to have been higher with lower exports



CDRR using ocean and Chipps Island recoveries versus just using Chipps Island recoveries (DRR) of the Mossdale or Durham Ferry and Jersey Point releases without the HORB in place.



No clear relationship between CDRR (DF/MD-JP survival) and flow/exports in years without a HORB.

What have we learned: No clear relationship between F/E ratio and smolt survival in range tested w/o barrier


Conceptual model of how smolt survival may vary with flow/exports

♦ no HORB

Escapement 2 1/2 years later

100000


Flow/export vs adult escapement




withHORB — Log. (no HORB)

 $R^2 = 0.5626 (p < 0.01)$ 80000 60000 40000 20000 0.1 10 100 1000


Mean Flow/Export Ratio between April 15 and June 15 (1951 to 2003) and escapement 2 ½ years later



VAMP release sites in the Sacramento-San Joaquin Estuary, California.



Results of ultrasonic tag detection (%) in upper Old River versus the San Joaquin River downstream of UOR during VAMP in 2006




Estimated SJ flow downstream of OR (in cfs)

Survival between Dos Reis and Jersey Point versus San Joaquin River flows downstream of Old River *.


Survival on the San Joaquin River generally higher with higher flows

^{*} Modeled DSM2 flows in 1990-2004, other estimate in 1989 and measured flows in 2005 and 2006



The CDRR for the Dos Reis group relative to the UOR group Those circled are significantly different than 1.0 at 95% CI

On average better survival for those that stay on San Joaquin River rather than migrate via UOR




DF MD DF DR DF DR MD DR MD

Combined Differential Recovery Rates (CDRR) (+ / - 2 SE) of CWT smolts released at Durham Ferry (DF), Mossdale (MD) and Dos Reis (DR) relative to those released at Jersey Point in 2000 – 2006.


What have we learned: Survival has been low since 2003

CDRR versus Vernalis flow with HORB



Survival for groups released at Mossdale and/or Durham Ferry relative to those released at Jersey Point and average flow at Vernalis in cfs

CDRR versus Vernalis flow with HORB



Relationship between flow and exports during VAMP tests with the HORB in place

What have we learned: Need to measure survival at exports of 1500 with a HORB in place

What should we do next?

- Measure exports of 1500 with 7000 flow with HORB
 - Continue testing with and without HORB
 - Identify sources/locations of mortality (Additional ultrasonic tagging work in south Delta proposed)